

IMO Shortlist 2005

Algebra

- 1 Find all pairs of integers a, b for which there exists a polynomial $P(x) \in \mathbb{Z}[X]$ such that product $(x^2 + ax + b) \cdot P(x)$ is a polynomial of a form

$$x^n + c_{n-1}x^{n-1} + \dots + c_1x + c_0$$

where each of c_0, c_1, \dots, c_{n-1} is equal to 1 or -1 .

- 2 We denote by \mathbb{R}^+ the set of all positive real numbers.

Find all functions $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ which have the property:

$$f(x)f(y) = 2f(x+yf(x))$$

for all positive real numbers x and y .

- 3 Four real numbers p, q, r, s satisfy $p + q + r + s = 9$ and $p^2 + q^2 + r^2 + s^2 = 21$. Prove that there exists a permutation (a, b, c, d) of (p, q, r, s) such that $ab - cd \geq 2$.
- 4 Find all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x+y) + f(x)f(y) = f(xy) + 2xy + 1$ for all real numbers x and y .
- 5 Let x, y, z be three positive reals such that $xyz \geq 1$. Prove that

$$\frac{x^5 - x^2}{x^5 + y^2 + z^2} + \frac{y^5 - y^2}{x^2 + y^5 + z^2} + \frac{z^5 - z^2}{x^2 + y^2 + z^5} \geq 0.$$

Hojoo Lee, Korea

Combinatorics

- 1] A house has an even number of lamps distributed among its rooms in such a way that there are at least three lamps in every room. Each lamp shares a switch with exactly one other lamp, not necessarily from the same room. Each change in the switch shared by two lamps changes their states simultaneously. Prove that for every initial state of the lamps there exists a sequence of changes in some of the switches at the end of which each room contains lamps which are on as well as lamps which are off.

- 2] This ISL 2005 problem has not been used in any TST I know. A pity, since it is a nice problem, but in its shortlist formulation, it is absolutely incomprehensible. Here is a mathematical restatement of the problem:

A forest consists of rooted (i. e. oriented) trees. Each vertex of the forest is either a leaf or has two successors. A vertex v is called an *extended successor* of a vertex u if there is a chain of vertices $u_0 = u, u_1, u_2, \dots, u_{t-1}, u_t = v$ such that the vertex u_{i+1} is a successor of the vertex u_i for every integer i with $0 \leq i \leq t-1$. A vertex is called *dynastic* if it has two successors and each of these successors has at least k extended successors.

Prove that if the forest has n vertices, then there are at most $\frac{n}{k+2}$ dynastic vertices.

- 3] Consider a $m \times n$ rectangular board consisting of mn unit squares. Two of its unit squares are called *adjacent* if they have a common edge, and a *path* is a sequence of unit squares in which any two consecutive squares are adjacent. Two paths are called *non-intersecting* if they don't share any common squares.

Each unit square of the rectangular board can be colored black or white. We speak of a *coloring* of the board if all its mn unit squares are colored.

Let N be the number of colorings of the board such that there exists at least one black path from the left edge of the board to its right edge. Let M be the number of colorings of the board for which there exist at least two non-intersecting black paths from the left edge of the board to its right edge.

Prove that $N^2 \geq M \cdot 2^{mn}$.

- 4] Let $n \geq 3$ be a fixed integer. Each side and each diagonal of a regular n -gon is labelled with a number from the set $\{1; 2; \dots; r\}$ in a way such that the following two conditions are fulfilled:
1. Each number from the set $\{1; 2; \dots; r\}$ occurs at least once as a label.
 2. In each triangle formed by three vertices of the n -gon, two of the sides are labelled with the same number, and this number is greater than the label of the third side.
- (a) Find the maximal r for which such a labelling is possible.

IMO Shortlist 2005

(b) *Harder version (IMO Shortlist 2005)*: For this maximal value of r , how many such labellings are there?

[hide="Easier version (5th German TST 2006) - contains answer to the harder version"]
Easier version (5th German TST 2006): Show that, for this maximal value of r , there are exactly $\frac{n!(n-1)!}{2^{n-1}}$ possible labellings.

- 5] There are n markers, each with one side white and the other side black. In the beginning, these n markers are aligned in a row so that their white sides are all up. In each step, if possible, we choose a marker whose white side is up (but not one of the outermost markers), remove it, and reverse the closest marker to the left of it and also reverse the closest marker to the right of it. Prove that, by a finite sequence of such steps, one can achieve a state with only two markers remaining if and only if $n - 1$ is not divisible by 3.
- 6] In a mathematical competition, in which 6 problems were posed to the participants, every two of these problems were solved by more than $\frac{2}{5}$ of the contestants. Moreover, no contestant solved all the 6 problems. Show that there are at least 2 contestants who solved exactly 5 problems each.

Radu Gologan and Dan Schwartz

- 7] Suppose that a_1, a_2, \dots, a_n are integers such that $n \mid a_1 + a_2 + \dots + a_n$. Prove that there exist two permutations (b_1, b_2, \dots, b_n) and (c_1, c_2, \dots, c_n) of $(1, 2, \dots, n)$ such that for each integer i with $1 \leq i \leq n$, we have

$$n \mid a_i - b_i - c_i$$

- 8] Suppose we have a n -gon. Some $n - 3$ diagonals are coloured black and some other $n - 3$ diagonals are coloured red (a side is not a diagonal), so that no two diagonals of the same colour can intersect strictly inside the polygon, although they can share a vertex. Find the maximum number of intersection points between diagonals coloured differently strictly inside the polygon, in terms of n .

IMO Shortlist 2005

Geometry

1] Given a triangle ABC satisfying $AC + BC = 3 \cdot AB$. The incircle of triangle ABC has center I and touches the sides BC and CA at the points D and E , respectively. Let K and L be the reflections of the points D and E with respect to I . Prove that the points A, B, K, L lie on one circle.

2] Six points are chosen on the sides of an equilateral triangle ABC : A_1, A_2 on BC , B_1, B_2 on CA and C_1, C_2 on AB , such that they are the vertices of a convex hexagon $A_1A_2B_1B_2C_1C_2$ with equal side lengths.

Prove that the lines A_1B_2, B_1C_2 and C_1A_2 are concurrent.

Bogdan Enescu, Romania

3] Let $ABCD$ be a parallelogram. A variable line g through the vertex A intersects the rays BC and DC at the points X and Y , respectively. Let K and L be the A -excenters of the triangles ABX and ADY . Show that the angle $\angle KCL$ is independent of the line g .

4] Let $ABCD$ be a fixed convex quadrilateral with $BC = DA$ and BC not parallel with DA . Let two variable points E and F lie of the sides BC and DA , respectively and satisfy $BE = DF$. The lines AC and BD meet at P , the lines BD and EF meet at Q , the lines EF and AC meet at R .

Prove that the circumcircles of the triangles PQR , as E and F vary, have a common point other than P .

5] Let $\triangle ABC$ be an acute-angled triangle with $AB \neq AC$. Let H be the orthocenter of triangle ABC , and let M be the midpoint of the side BC . Let D be a point on the side AB and E a point on the side AC such that $AE = AD$ and the points D, H, E are on the same line. Prove that the line HM is perpendicular to the common chord of the circumscribed circles of triangle $\triangle ABC$ and triangle $\triangle ADE$.

[url=<http://www.mathlinks.ro/Forum/viewtopic.php?t=89094>]**All the problems from the Swiss Imo S**

6] Let ABC be a triangle, and M the midpoint of its side BC . Let γ be the incircle of triangle ABC . The median AM of triangle ABC intersects the incircle γ at two points K and L . Let the lines passing through K and L , parallel to BC , intersect the incircle γ again in two points X and Y . Let the lines AX and AY intersect BC again at the points P and Q . Prove that $BP = CQ$.

7] In an acute triangle ABC , let D, E, F be the feet of the perpendiculars from the points A, B, C to the lines BC, CA, AB , respectively, and let P, Q, R be the feet of the perpendiculars from the points A, B, C to the lines EF, FD, DE , respectively.

IMO Shortlist 2005

Prove that $p(ABC)p(PQR) \geq (p(DEF))^2$, where $p(T)$ denotes the perimeter of triangle T .

Number Theory

- [1] Determine all positive integers relatively prime to all the terms of the infinite sequence

$$a_n = 2^n + 3^n + 6^n - 1, \quad n \geq 1.$$

- [2] Let a_1, a_2, \dots be a sequence of integers with infinitely many positive and negative terms. Suppose that for every positive integer n the numbers a_1, a_2, \dots, a_n leave n different remainders upon division by n .

Prove that every integer occurs exactly once in the sequence a_1, a_2, \dots

- [3] Let a, b, c, d, e, f be positive integers and let $S = a + b + c + d + e + f$. Suppose that the number S divides $abc + def$ and $ab + bc + ca - de - ef - df$. Prove that S is composite.

- [4] Find all positive integers n such that there exists a unique integer a such that $0 \leq a < n!$ with the following property:

$$n! \mid a^n + 1$$

- [5] Denote by $d(n)$ the number of divisors of the positive integer n . A positive integer n is called highly divisible if $d(n) > d(m)$ for all positive integers $m < n$. Two highly divisible integers m and n with $m < n$ are called consecutive if there exists no highly divisible integer s satisfying $m < s < n$.

(a) Show that there are only finitely many pairs of consecutive highly divisible integers of the form (a, b) with $a \mid b$.

(b) Show that for every prime number p there exist infinitely many positive highly divisible integers r such that pr is also highly divisible.

- [6] Let a, b be positive integers such that $b^n + n$ is a multiple of $a^n + n$ for all positive integers n . Prove that $a = b$.

- [7] Let $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$, where a_0, \dots, a_n are integers, $a_n > 0$, $n \geq 2$. Prove that there exists a positive integer m such that $P(m!)$ is a composite number.