

## Algebra

- 1  $a_0, a_1, a_2, \dots$  is a sequence of real numbers such that

$$a_{n+1} = [a_n] \cdot \{a_n\}$$

prove that exist  $j$  such that for every  $i \geq j$  we have  $a_{i+2} = a_i$ .

- 2 Let  $a_0, a_1, a_2, \dots$  be a sequence of reals such that  $a_0 = -1$  and

$$a_n + \frac{a_{n-1}}{2} + \frac{a_{n-2}}{3} + \dots + \frac{a_1}{n} + \frac{a_0}{n+1} = 0 \text{ for all } n \geq 1.$$

Show that  $a_n > 0$  for all  $n \geq 1$ .

- 3 The sequence  $c_0, c_1, \dots, c_n, \dots$  is defined by  $c_0 = 1, c_1 = 0$ , and  $c_{n+2} = c_{n+1} + c_n$  for  $n \geq 0$ . Consider the set  $S$  of ordered pairs  $(x, y)$  for which there is a finite set  $J$  of positive integers such that  $x = \sum_{j \in J} c_j, y = \sum_{j \in J} c_{j-1}$ . Prove that there exist real numbers  $\alpha, \beta$ , and  $M$  with the following property: An ordered pair of nonnegative integers  $(x, y)$  satisfies the inequality  $m < \alpha x + \beta y < M$  if and only if  $(x, y) \in S$ .

*Remark:* A sum over the elements of the empty set is assumed to be 0.

- 4 Prove the inequality:

$$\sum_{i < j} \frac{a_i a_j}{a_i + a_j} \leq \frac{n}{2(a_1 + a_2 + \dots + a_n)} \cdot \sum_{i < j} a_i a_j$$

for positive reals  $a_1, a_2, \dots, a_n$ .

- 5 If  $a, b, c$  are the sides of a triangle, prove that

$$\sum_{\text{cyc}} \frac{\sqrt{a+b-c}}{\sqrt{a} + \sqrt{b} - \sqrt{c}} \leq 3$$

- 6 Determine the least real number  $M$  such that the inequality  $|ab(a^2 - b^2) + bc(b^2 - c^2) + ca(c^2 - a^2)| \leq M$  holds for all real numbers  $a, b$  and  $c$ .

## Combinatorics

- 1 We have  $n \geq 2$  lamps  $L_1, \dots, L_n$  in a row, each of them being either on or off. Every second we simultaneously modify the state of each lamp as follows: if the lamp  $L_i$  and its neighbours (only one neighbour for  $i = 1$  or  $i = n$ , two neighbours for other  $i$ ) are in the same state, then  $L_i$  is switched off; otherwise,  $L_i$  is switched on. Initially all the lamps are off except the leftmost one which is on.

(a) Prove that there are infinitely many integers  $n$  for which all the lamps will eventually be off. (b) Prove that there are infinitely many integers  $n$  for which the lamps will never be all off.

- 2 Let  $P$  be a regular 2006-gon. A diagonal is called *good* if its endpoints divide the boundary of  $P$  into two parts, each composed of an odd number of sides of  $P$ . The sides of  $P$  are also called *good*. Suppose  $P$  has been dissected into triangles by 2003 diagonals, no two of which have a common point in the interior of  $P$ . Find the maximum number of isosceles triangles having two good sides that could appear in such a configuration.

- 3 Let  $S$  be a finite set of points in the plane such that no three of them are on a line. For each convex polygon  $P$  whose vertices are in  $S$ , let  $a(P)$  be the number of vertices of  $P$ , and let  $b(P)$  be the number of points of  $S$  which are outside  $P$ . A line segment, a point, and the empty set are considered as convex polygons of 2, 1, and 0 vertices respectively. Prove that for every real number  $x$ :

$$\sum_P x^{a(P)}(1-x)^{b(P)} = 1, \text{ where the sum is taken over all convex polygons with vertices in } S.$$

*Alternative formulation:*

Let  $M$  be a finite point set in the plane and no three points are collinear. A subset  $A$  of  $M$  will be called *round* if its elements is the set of vertices of a convex  $A$ -gon  $V(A)$ . For each round subset let  $r(A)$  be the number of points from  $M$  which are exterior from the convex  $A$ -gon  $V(A)$ . Subsets with 0, 1 and 2 elements are always round, its corresponding polygons are the empty set, a point or a segment, respectively (for which all other points that are not vertices of the polygon are exterior). For each round subset  $A$  of  $M$  construct the polynomial

$$P_A(x) = x^{|A|}(1-x)^{r(A)}.$$

Show that the sum of polynomials for all round subsets is exactly the polynomial  $P(x) = 1$ .

- 4 A cake has the form of an  $n \times n$  square composed of  $n^2$  unit squares. Strawberries lie on some of the unit squares so that each row or column contains exactly one strawberry; call this arrangement  $A$ . Let  $B$  be another such arrangement. Suppose that every grid rectangle with

# IMO Shortlist 2006

---

one vertex at the top left corner of the cake contains no fewer strawberries of arrangement  $B$  than of arrangement  $A$ .

Prove that arrangement  $B$  can be obtained from  $A$  by performing a number of switches, defined as follows: A switch consists in selecting a grid rectangle with only two strawberries, situated at its top right corner and bottom left corner, and moving these two strawberries to the other two corners of that rectangle.

- 5 An  $(n, k)$ -tournament is a contest with  $n$  players held in  $k$  rounds such that:
- (i) Each player plays in each round, and every two players meet at most once.
  - (ii) If player  $A$  meets player  $B$  in round  $i$ , player  $C$  meets player  $D$  in round  $i$ , and player  $A$  meets player  $C$  in round  $j$ , then player  $B$  meets player  $D$  in round  $j$ .

Determine all pairs  $(n, k)$  for which there exists an  $(n, k)$ -tournament.

- 6 A holey triangle is an upward equilateral triangle of side length  $n$  with  $n$  upward unit triangular holes cut out. A diamond is a  $60^\circ - 120^\circ$  unit rhombus. Prove that a holey triangle  $T$  can be tiled with diamonds if and only if the following condition holds: Every upward equilateral triangle of side length  $k$  in  $T$  contains at most  $k$  holes, for  $1 \leq k \leq n$ .

- 7 Consider a convex polyhedron without parallel edges and without an edge parallel to any face other than the two faces adjacent to it. Call a pair of points of the polyhedron *antipodal* if there exist two parallel planes passing through these points and such that the polyhedron is contained between these planes. Let  $A$  be the number of antipodal pairs of vertices, and let  $B$  be the number of antipodal pairs of midpoint edges. Determine the difference  $A - B$  in terms of the numbers of vertices, edges, and faces.

# IMO Shortlist 2006

---

## Geometry

- 1] Let  $ABC$  be triangle with incenter  $I$ . A point  $P$  in the interior of the triangle satisfies

$$\angle PBA + \angle PCA = \angle PBC + \angle PCB.$$

Show that  $AP \geq AI$ , and that equality holds if and only if  $P = I$ .

- 2] Let  $ABCD$  be a trapezoid with parallel sides  $AB > CD$ . Points  $K$  and  $L$  lie on the line segments  $AB$  and  $CD$ , respectively, so that  $\frac{AK}{KB} = \frac{DL}{LC}$ . Suppose that there are points  $P$  and  $Q$  on the line segment  $KL$  satisfying  $\angle APB = \angle BCD$  and  $\angle CQD = \angle ABC$ . Prove that the points  $P, Q, B$  and  $C$  are concyclic.

- 3] Consider a convex pentagon  $ABCDE$  such that

$$\angle BAC = \angle CAD = \angle DAE \quad , \quad \angle ABC = \angle ACD = \angle ADE$$

Let  $P$  be the point of intersection of the lines  $BD$  and  $CE$ . Prove that the line  $AP$  passes through the midpoint of the side  $CD$ .

- 4] Let  $ABC$  be a triangle such that  $\widehat{ACB} < \widehat{BAC} < \frac{\pi}{2}$ . Let  $D$  be a point of  $[AC]$  such that  $BD = BA$ . The incircle of  $ABC$  touches  $[AB]$  at  $K$  and  $[AC]$  at  $L$ . Let  $J$  be the center of the incircle of  $BCD$ . Prove that  $(KL)$  intersects  $[AJ]$  at its middle.

- 5] In triangle  $ABC$ , let  $J$  be the center of the excircle tangent to side  $BC$  at  $A_1$  and to the extensions of the sides  $AC$  and  $AB$  at  $B_1$  and  $C_1$  respectively. Suppose that the lines  $A_1B_1$  and  $AB$  are perpendicular and intersect at  $D$ . Let  $E$  be the foot of the perpendicular from  $C_1$  to line  $DJ$ . Determine the angles  $\angle BEA_1$  and  $\angle AEB_1$ .

- 6] Circles  $w_1$  and  $w_2$  with centres  $O_1$  and  $O_2$  are externally tangent at point  $D$  and internally tangent to a circle  $w$  at points  $E$  and  $F$  respectively. Line  $t$  is the common tangent of  $w_1$  and  $w_2$  at  $D$ . Let  $AB$  be the diameter of  $w$  perpendicular to  $t$ , so that  $A, E, O_1$  are on the same side of  $t$ . Prove that lines  $AO_1, BO_2, EF$  and  $t$  are concurrent.

- 7] In a triangle  $ABC$ , let  $M_a, M_b, M_c$  be the midpoints of the sides  $BC, CA, AB$ , respectively, and  $T_a, T_b, T_c$  be the midpoints of the arcs  $BC, CA, AB$  of the circumcircle of  $ABC$ , not containing the vertices  $A, B, C$ , respectively. For  $i \in \{a, b, c\}$ , let  $w_i$  be the circle with  $M_iT_i$  as diameter. Let  $p_i$  be the common external common tangent to the circles  $w_j$  and  $w_k$  (for all  $\{i, j, k\} = \{a, b, c\}$ ) such that  $w_i$  lies on the opposite side of  $p_i$  than  $w_j$  and  $w_k$  do. Prove that the lines  $p_a, p_b, p_c$  form a triangle similar to  $ABC$  and find the ratio of similitude.

# IMO Shortlist 2006

---

- 8] Let  $ABCD$  be a convex quadrilateral. A circle passing through the points  $A$  and  $D$  and a circle passing through the points  $B$  and  $C$  are externally tangent at a point  $P$  inside the quadrilateral. Suppose that  $\angle PAB + \angle PDC \leq 90^\circ$  and  $\angle PBA + \angle PCD \leq 90^\circ$ . Prove that  $AB + CD \geq BC + AD$ .
- 9] Points  $A_1, B_1, C_1$  are chosen on the sides  $BC, CA, AB$  of a triangle  $ABC$  respectively. The circumcircles of triangles  $AB_1C_1, BC_1A_1, CA_1B_1$  intersect the circumcircle of triangle  $ABC$  again at points  $A_2, B_2, C_2$  respectively ( $A_2 \neq A, B_2 \neq B, C_2 \neq C$ ). Points  $A_3, B_3, C_3$  are symmetric to  $A_1, B_1, C_1$  with respect to the midpoints of the sides  $BC, CA, AB$  respectively. Prove that the triangles  $A_2B_2C_2$  and  $A_3B_3C_3$  are similar.
- [hide="Comment"]This is my personal favourite of the ISL Geometry problems ¡!- s:D - ;img src="SMILIES\_PATH/icon\_mrgreen.gif" alt = " : D" title = "Mr.Green" / ><! - - s : D - - >
- 10] Assign to each side  $b$  of a convex polygon  $P$  the maximum area of a triangle that has  $b$  as a side and is contained in  $P$ . Show that the sum of the areas assigned to the sides of  $P$  is at least twice the area of  $P$ .

## Number Theory

- 1 Determine all pairs  $(x, y)$  of integers such that

$$1 + 2^x + 2^{2x+1} = y^2.$$

- 2 For  $x \in (0, 1)$  let  $y \in (0, 1)$  be the number whose  $n$ -th digit after the decimal point is the  $2^n$ -th digit after the decimal point of  $x$ . Show that if  $x$  is rational then so is  $y$ .

- 3 We define a sequence  $(a_1, a_2, a_3, \dots)$  by setting

$$a_n = \frac{1}{n} \left( \left[ \frac{n}{1} \right] + \left[ \frac{n}{2} \right] + \dots + \left[ \frac{n}{n} \right] \right)$$

for every positive integer  $n$ . Hereby, for every real  $x$ , we denote by  $[x]$  the integral part of  $x$  (this is the greatest integer which is  $\leq x$ ).

**a)** Prove that there is an infinite number of positive integers  $n$  such that  $a_{n+1} > a_n$ . **b)** Prove that there is an infinite number of positive integers  $n$  such that  $a_{n+1} < a_n$ .

- 4 Let  $P(x)$  be a polynomial of degree  $n > 1$  with integer coefficients and let  $k$  be a positive integer. Consider the polynomial  $Q(x) = P(P(\dots P(P(x)) \dots))$ , where  $P$  occurs  $k$  times. Prove that there are at most  $n$  integers  $t$  such that  $Q(t) = t$ .

- 5 Prove that the equation  $\frac{x^7-1}{x-1} = y^5 - 1$  doesn't have integer solutions!

- 6 Let  $a > b > 1$  be relatively prime positive integers. Define the weight of an integer  $c$ , denoted by  $w(c)$  to be the minimal possible value of  $|x| + |y|$  taken over all pairs of integers  $x$  and  $y$  such that  $ax + by = c$ . An integer  $c$  is called a *local champion* if  $w(c) \geq w(c \pm a)$  and  $w(c) \geq w(c \pm b)$ . Find all local champions and determine their number.

- 7 For all positive integers  $n$ , show that there exists a positive integer  $m$  such that  $n$  divides  $2^m + m$ .